


Most people have successfully lost weight at some point in their lives. However, people's success at maintaining that weight loss is another story. When you look at clinical trials of long-term weight loss maintenance, the numbers don't look so good. [Approximately 10-20% of subjects are able to maintain a weight loss of at least 5% after 5 years.](#) The results of other research also look grim; [five years after completing structured weight loss programs, people only maintain a 3% weight loss.](#)

The problem with clinical trials, however, is that they may not reflect what is going on with the typical U.S. citizen. [A recent study](#) used data from the National Health and Nutrition Examination (NHANES) survey to get an idea of the success rates of the average person in the U.S. The researchers looked at data for 14,306 people between 20 and 84 years of age, from the years 1999 to 2006. They defined "long-term weight loss maintenance" as having maintained weight loss for at least 1 year. They excluded people who had never been overweight or obese.

The following chart shows the results of the study:



You can see that over 35% of people were successful at maintaining a weight loss of at least 5% for more than a year. On average, this was a 29 pound weight loss. The numbers drop off from there, with only 17% maintaining a 10% weight loss. However, of the people in that 17%, the average weight weight loss maintained was 42 pounds.

Study Shortcomings

The results of this study suggest that Americans are doing a better job of maintaining weight loss than clinical trials would have you believe. However, there are a number of limitations to this study that you need to consider:

- The research involved self report of body weight, which could introduce bias because [people tend to under-report their weight](#). Also, since this data was based on individuals trying to remember their past weights, there could be recall bias. However, there is some research to indicate that, on the whole, [recall of past weight is reasonable stable over time](#). So recall bias probably does not affect the results of this study too much.
- The researchers were not able to determine the method of weight loss. Thus, the data could include people who had unintentional weight loss due to illness. Since this was a cross-sectional study, the researchers could not determine when the weight loss occurred. The researchers also could not determine whether the people who had large amounts of weight loss underwent bariatric surgery; however, the number of people who have such surgery is very small (only 39 per 100,000 people), so this probably did not affect the results.
- The biggest limitation of this study is that the researchers only looked at one point in time. They only had data on the people's current weights, maximum weights, and weight 1 year ago. That's a lot of missing years that remain unaccounted. Thus, there was no way to determine the number of years people had maintained their weight loss. Still, the study does give at least some idea of long-term weight loss success. For example, someone who had a maximum weight of 220 pounds, went down 165 pounds two years ago, and then went to 185 pounds, would be classified as having 10% long-term weight maintenance in this study. While the person certainly regained some weight, he/she didn't regain all of the weight, and thus are still more successful at weight loss than if he/she had gone back to a maximum weight.

Better Than Clinical Trials, But Still Bad

Overall, the data does suggest that may do a bit better at maintaining long-term weight loss than clinical trials indicate. Still, if only 17% are maintaining a 10% long-term weight loss, that means 83% of people are regaining most, if not all, of their weight back. Those aren't good odds.

While the statistics are grim, it doesn't mean that **you** can't be successful at maintaining long-term weight loss. But long-term weight loss requires effort, and an effective strategy. If you are someone who has struggled with maintaining weight loss, you can [contact me for online consultation](#). I am also available to [speak to groups and organizations on effective strategies for maintaining long term weight loss](#). With appropriate lifestyle modifications, you

can be in that 17%.

REFERENCE: [Kraschnewski, J.L., et al. Long-term weight loss maintenance in the United States. International Journal of Obesity \(published ahead of print; available online May 18, 2010\)](#)

[CLICK HERE to find out more about the Weightology RESEARCH REVIEW.](#)

[CLICK HERE to learn about ONLINE COACHING.](#)