


[A recent study](#) out of Princeton University has the [high-fructose corn syrup alarmists](#) out in full force. This study compared the effects of high fructose corn syrup (HFCS) to regular table sugar (sucrose), looking at their effects on body weight, body fat, and triglycerides (fats that float around in your blood). The study found that the rats fed HFCS gained more weight and abdominal fat than the rats fed sucrose. This study has strengthened the belief of some people that HFCS is contributing to obesity in our society, and that it is worse than regular sugar. But is it really?

To answer this question, we need to take a close look at this study. The researchers performed 2 experiments. In the first experiment, male rats were divided into 4 groups. Group 1 (the control group) was fed a regular diet. Group 2 was fed the same diet, with the addition of 24-hour access to water sweetened with HFCS. Group 3 had the regular diet with 12-hour access to the HFCS-sweetened water. Group 4 had the regular diet, with 12-hour access to sucrose-sweetened water. The rats were tracked for 8 weeks; weight was measured, along with food, sucrose, and HFCS intake.

You can see the results for experiment 1 in the following chart:




The rats who got HFCS for 12 hours gained significantly more weight than the other 3 groups. At first glance, this would make you believe that HFCS makes you gain more weight than sucrose, even if you are eating the same number of calories. However, there is a problem with these results. Take a look again at the chart above. If the rats fed HFCS for 12 hours gained more weight, why didn't the rats fed HFCS for 24 hours also gain more weight? They got HFCS for a full 12 hours more, yet didn't gain more weight. This is a glaring inconsistency in the results...an inconsistency that the researchers never tried to explain.

Rather than some unique effect of HFCS, a more likely explanation is one of chance. Put on your math hat, because we need to talk about some statistics. Researchers use statistics to get an idea of the probability that their results are due to chance. When the scientists run their stats, they get what is known as a P value. The P value tells you the probability that the results are not due to chance. Usually, if the P value is less than 0.05, a scientist will call the results “significant.” In other words, if you did the experiment 100 times, you would only see these results less than 5 times if there wasn’t a true effect.

The above only holds true if you’re doing a single comparison. If you start comparing a bunch of groups all to each other, the probability of a fluke result dramatically increases. The Princeton study is a perfect example. There are 4 groups all being compared to each other. That makes for 6 total comparisons (group 1 to group 2, 1 to 3, 1 to 4, 2 to 3, 2 to 4, and 3 to 4). Each one of these comparisons is being tested against that 5% level. To calculate the probability of a fluke result in this case, we calculate $1 - (0.95 \times 0.95 \times 0.95 \times 0.95 \times 0.95 \times 0.95) = 26\%$. In other words, there is a 1 in 4 chance that the greater weight gain in the HFCS-fed rats is a fluke. I don’t know about you, but I wouldn’t put too much faith in results that have a 1 in 4 chance of being wrong. There are ways that scientists can adjust for this, but the Princeton researchers didn’t appear to make those adjustments. Thus, it is not surprising that there was a significant result observed in 1 out of the 4 groups...you would expect this to happen based on random chance alone.

In Experiment 2, the researchers divided male rats into 3 groups: 12-hour HFCS, 24-hour HFCS, and control. They tracked the rats for 6 months. Both HFCS-fed groups gained more weight and fat than the control, and also had higher triglycerides. However, the researchers didn’t compare HFCS to sucrose in this group, so this experiment doesn’t say anything about whether HFCS is any worse than sucrose. The researchers also didn’t say anything about food intake and whether the HFCS-fed rats ate more than the control rats.

Experiment 2 also featured female rats on one of the 4 diets used in Experiment 1. These rats were tracked for 7 months. The following chart shows the results of the experiment:



The female rats fed HFCS for 24 hours a day gained significantly more weight than the other groups. Now compare these results to the chart for Experiment 1 earlier. Do you see the disparity? In Experiment 1, the rats fed HFCS for 12 hours per day gained the most weight. However, in Experiment 2, the rats fed HFCS for 24 hours per day gained the most weight, and the female rats fed HFCS for 12 hours didn't gain any more weight than the other groups. Why did the 12-hour group gain the most weight in one experiment, but the 24-hour group gain the most weight in a nearly identical experiment? This is a glaring contradiction in the results, and a problem which the researchers did not discuss. We also have the same statistical problem that we did with Experiment 1. Since there are 6 comparisons, there is a 1 in 4 chance that the results are wrong (and ironically, we have 1 out of the 4 groups showing a significant result). In fact, when we take both experiments combined, we have at least a 50% chance that the results of one of the experiments are wrong. Out of all the comparisons being made, we would expect to see a couple groups show a significant result based on random chance...and that's exactly what happened in this study.

The bottom line is that there is no valid reason for HFCS to be any different than sucrose in the way that it affects your body. They are both nearly identical in their composition, containing roughly half fructose and half glucose. They are both nearly identical in the way they are metabolized by your body. There is no practical difference between the two as far as your body is concerned. Now, I'm not saying that you should go out and consume all the HFCS that you want. The point is that there is nothing uniquely "bad" about HFCS compared to regular sugar. HFCS is not uniquely responsible for weight gain as some people would have you believe.

If you see a product with HFCS and a similar product with natural table sugar, don't assume the product with natural sugar is any better. Rather than worrying about whether something

contains HFCS, you should strive to reduce your intake of all types of added sugar and refined carbohydrates in your diet. It is much more important to look at the big picture; keep your physical activity high, manage your overall food intake, make sure most of your food is from minimally refined sources, and keep your protein intake high. This is what will help you lose weight and keep it off, rather than singling out HFCS in your diet. Don't let the fructose fear-mongers fool you.

REFERENCE: [Bocarsly, M.E., et al. High-fructose corn syrup causes characteristics of obesity in rats: Increased body weight, body fat and triglyceride levels. *Pharmacology, Biochemistry and Behavior* \(published ahead of print; available online February 26, 2010\)](#)